


RD-A142 783 MACROCYCLES CONTAINING TIN SYNTHESIS AND STRUCTURE OF
110-DIPHENYL-110-DI. (U) TEXAS A AND M UNIV COLLEGE
STATION DEPT OF CHEMISTRY M NEWCOMB ET AL. 02 JUL 84
UNCLASSIFIED N80014-79-C-0584 F/G 7/3 NL





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

AD-A142 703

DTIC FILE COPY

Unclassified

12

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER Technical Report 5	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Macrocycles Containing Tin. Synthesis and Structure of 1,10-Diphenyl-1,10-distanna-bicycle[8.8.8]hexacosane		5. TYPE OF REPORT & PERIOD COVERED Technical Report
7. AUTHOR(s) Martin Newcomb, Michael T. Blanda, Yutaka Azuma, Terry J. Delord		6. PERFORMING ORG. REPORT NUMBER N00014-79-C-0584
9. PERFORMING ORGANIZATION NAME AND ADDRESS Texas A&M University Department of Chemistry College Station, TX 77843		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 053-714
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Department of the Navy Arlington, Virginia 22217		12. REPORT DATE July 2, 1984
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		13. NUMBER OF PAGES 11
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES To be published in <u>J. Chem. Soc., Chem. Commun.</u>		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Macrocycle X-ray crystallography Macrobicycle Structure Polystannane Synthesis		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Synthetic methods used for the preparation and purification of the title compound, a representative member of a new class of compounds, are reported. The macrobicycle employs tin atoms as bridgeheads in a manner analogous to the nitrogen atoms of a cryptand. The molecular and crystal structure of the macrobicycle were determined by x-ray crystallography, and the positional and thermal parameters and bond distances are reported.		

DD FORM 1 JAN 73 1473 EDITION OF 1 NOV 68 IS OBSOLETE
S/N 0102-014-6601

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)



OFFICE OF NAVAL RESEARCH

Contract N00014-79-C-0584

Task No. NR 053-714

TECHNICAL REPORT NO. 5

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Avail and/or	
Dist	Special

A-1

Macrocycles Containing Tin.

Synthesis and Structure of

1,10-Diphenyl-1,10-distannabicyclo[8.8.8]hexacosane

by

Martin Newcomb, Michael T. Blanda, Yutaka Azuma, and Terry J. Delord

Texas A&M University
Department of Chemistry
College Station, TX 77843

July 2, 1984

Prepared for publication in J. Chem. Soc., Chem. Commun.Reproduction in whole or in part is permitted for
any purpose of the United States Government* This document has been approved for public release
and sale; its distribution is unlimited* This statement should also appear in Item 16 of Document Control Data
- DD Form 1473. Copies of form available from cognizant contract
administrator.

84 07 05 076

Macrocycles Containing Tin. Synthesis and Structure of
1,10-Diphenyl-1,10-distannabicyclo[8.8.8]hexacosane

Martin Newcomb*,†, Michael T. Blanda, Yutaka Azuma, and Terry J. Delord
Department of Chemistry, Texas A&M University, College Station, TX, 77843 USA

Summary: The title macrobicycle was prepared and studied by x-ray crystallography.

We have reported the syntheses of macrocycles containing two or four diphenylstanna moieties in the ring with various length polymethylene chains between the tin atoms.¹ Since acyclic, Lewis acidic distanna compounds have been shown to exhibit cooperative binding of donors,² we predicted that macrocycles containing Lewis acidic tin atoms would display cooperative and size-selective binding properties. In confirmation of this expectation, we have found that replacement of one or both phenyl groups on each tin atom in a macrocycle with chlorine or bromine atoms gives Lewis acidic macrocycles which form complexes with donor compounds containing basic nitrogen and oxygen atoms; complexes with donor atom to tin atom ratios of both 2:1 and 1:1 have been isolated and are under investigation.³ In this communication we report the preparation and structure of a representative of a new class of organotin compounds, a macrobicycle containing two tin atoms in which the tin atoms serve as bridgeheads in a manner analogous to the nitrogen atoms of a cryptand. It may be expected that Lewis acidic macrobicyclic organotin compounds will be highly selective complexing agents.

The preparation of 1,10-diphenyl-1,10-distannabicyclo[8.8.8]hexacosane (**1**) follows the sequence shown in the Scheme. The preparation of macrocycle **4** has been reported,¹ but we modified the procedure by using a double dilution method wherein dilute (ca. 0.03 M) solutions of the reagents **2** and **3**¹ were added

simultaneously to a THF reaction mixture over 2 h at 25 °C to give **4** in 62% yield after purification by reverse phase chromatography¹ and recrystallization. Treatment of macrocycle **4** with 2.1 molar equivalents of HCl in CH₂Cl₂ at -78 °C followed by slow warming to 25 °C and distillation of the solvent at reduced pressure gave the dichlorinated product **5** which appeared to be pure by ¹H NMR spectroscopy. Crude **5** was then allowed to react with the diGrignard reagent **2** in THF at 25 °C, again using the double dilution method (0.03 M solutions, 2 h addition). After a standard workup,¹ bicyclic **1** was purified by preparative reverse phase chromatography (Baker bonded-phase ODS, 40 μ; 60:40, CH₃OH:THF elution; ca. 5 column volume retention) followed by recrystallization from CH₃OH/THF to give pure **1** in 15% yield from macrocycle **4**. Macrobicyclic **1** (mp 92–94 °C) was characterized by ¹H NMR and ¹³C NMR spectroscopy, osmometric MW measurement (Calcd: 728; found: 734), and elemental analysis (C, H).

A crystal of **1** was suitable for x-ray crystallographic analysis.[†] An ORTEP drawing of **1** is shown in Figure 1. The compound crystallized in a cylindrical form with a tin-tin distance of 8.45 Å. The interior of the cavity in **1** is filled with the hydrogen atoms of the polymethylene chains. The tin atoms of **1** are essentially tetrahedral in the crystal, and from the ¹J(¹¹⁹Sn–¹³C) of 332 Hz appear to be tetrahedral in solution.⁴

Crystal data: Sn₂C₃₆H₅₈, M = 728.24, triclinic, space group P1, a = 13.508(8), b = 16.154(9), c = 9.094(3) Å, α = 106.35(4)°, β = 109.67(4)°, γ = 92.49(5)°, U = 1772(1) Å³, Z = 2, D_c = 1.365 g/cm³. Intensity data were collected on an Enraf-Nonius CAD-4 diffractometer with Mo-Kα radiation. Data were corrected for Lorentz, polarization and absorption effects (μ = 14.38 cm⁻¹) to yield 2193 observed structure factors with I > 3σ(I) which were used for the structure solution and refinement. The positions of the two tin atoms were determined by direct methods. Successive least squares refinement and difference-Fourier methods were used to

locate the remaining carbon atoms. While exhibiting rather high thermal motion, all non-hydrogen atoms were successfully refined anisotropically. The hydrogen atoms were included in calculated positions with fixed temperature factors and were not refined. Least squares refinement converged at $R = 0.061$ and $R_w = 0.084$.

Other size distanna macrobicycles should be available by the procedures we used to prepare **1**, and, indeed, the corresponding 32 atom bicycle has already been prepared in our laboratory. In addition, functionalization of macrobicycle **1** to give a Lewis acidic species (replacement of phenyl groups with chloro groups) was achieved by treatment of **1** with HCl in a reaction similar to that used for conversion of **4** to **5**, and complexation properties of this derivative are under investigation.³

We are grateful for financial support from the Office of Naval Research.

References

1. Y. Azuma and M. Newcomb, Organometallics, 1984, **3**, 9.
2. (a) T. J. Karol, J. P. Hutchinson, J. R. Hyde, H. G. Kuivila, and J. A. Zubieta, Organometallics, 1983, **2**, 106. (b) K. Jurkschat and M. Gielen, Bull. Soc. Chim. Belg., 1982, **91**, 803.
3. M. Newcomb, M. T. Blanda, R. M. Sanchez, Y. Azuma; unpublished results.
4. H. G. Kuivila, J. L. Considine, R. H. Sarma, R. J. Mynott, J. Organometal. Chem., 1976, **111**, 179.

[†]Camille and Henry Dreyfus Teacher-Scholar, 1980-1985.

[‡]All crystallographic computing and plotting was performed using the Enraf-Nonius Structure Determination Package (1982), Enraf-Nonius, Delft, Holland. The atomic coordinates for this work are available on request from the Director of the Cambridge Crystallographic Data Centre, University Chemical Laboratory, Lensfield Rd., Cambridge CB2 1EW. Any request should be accompanied by the full literature citation for this communication.




Figure 1. ORTEP representation of **1**. The hydrogen atoms have been omitted for clarity.

Scheme




Table of Positional and Thermal Parameters and Their Estimated Standard Deviations.

Atom	x	y	z	<i>A</i> (1, 1)	<i>A</i> (2, 2)	<i>A</i> (3, 3)	<i>A</i> (1, 2)	<i>A</i> (1, 3)	<i>A</i> (2, 3)
SN1	0.32226(9)	0.52545(6)	0.1821(1)	0.00689(1)	0.0316(5)	0.0083(1)	0.0167(2)	0.0096(2)	0.0138(1)
SN2	0.1781(8)	-0.02537(6)	-0.1653(1)	0.0182(1)	0.00700(5)	0.0392(2)	0.0006(1)	0.0273(2)	0.017(2)
C1	0.339(1)	0.6657(8)	0.259(1)	0.014(1)	0.0098(7)	0.038(2)	0.010(1)	0.020(2)	0.017(2)
C2	0.293(1)	0.7121(9)	0.359(1)	0.018(1)	0.0093(8)	0.038(2)	0.004(2)	0.032(2)	0.017(2)
C3	0.356(1)	0.8524(9)	0.485(2)	0.017(1)	0.0059(7)	0.055(4)	0.002(2)	0.026(3)	0.017(3)
C4	0.367(1)	0.8494(8)	0.363(2)	0.017(2)	0.0048(6)	0.061(4)	0.003(2)	0.019(4)	0.012(2)
C5	0.412(1)	0.8843(8)	0.263(2)	0.014(1)	0.0091(7)	0.054(3)	0.004(2)	0.018(3)	0.026(2)
C6	0.454(1)	0.7141(9)	0.219(1)	0.019(1)	0.0105(8)	0.029(2)	0.013(2)	0.027(2)	0.014(2)
C7	0.162(1)	-0.1632(8)	-0.245(1)	0.013(1)	0.0058(6)	0.034(2)	0.005(1)	0.013(3)	0.006(2)
C8	0.596(1)	-0.2136(8)	-0.457(1)	0.015(1)	0.0095(7)	0.037(2)	0.012(2)	0.025(3)	0.019(2)
C9	0.587(1)	-0.3828(9)	-0.458(2)	0.014(1)	0.0091(9)	0.037(2)	0.012(2)	0.029(3)	0.019(2)
C10	0.133(1)	-0.3459(9)	-0.355(2)	0.017(1)	0.0072(8)	0.041(3)	0.004(2)	0.028(3)	0.008(3)
C11	0.193(1)	-0.3887(8)	-0.281(2)	0.023(2)	0.0066(6)	0.043(3)	0.006(2)	0.015(4)	0.014(2)
C12	0.218(1)	-0.2189(9)	-0.147(2)	0.016(2)	0.0077(8)	0.030(3)	0.006(2)	0.009(3)	0.003(2)
C13	0.256(2)	0.4763(12)	-0.577(2)	0.021(2)	0.0153(12)	0.054(4)	0.027(2)	0.032(3)	0.022(3)
C14	0.265(2)	0.4848(15)	-0.178(2)	0.021(2)	0.0254(20)	0.035(4)	0.023(3)	0.014(4)	0.007(4)
C15	0.225(1)	0.3698(11)	-0.353(2)	0.017(1)	0.0125(11)	0.029(3)	0.006(2)	0.013(3)	0.008(3)
C16	0.289(1)	0.2717(11)	-0.424(2)	0.028(2)	0.0145(11)	0.038(3)	0.015(3)	0.033(4)	0.022(3)
C17	0.294(1)	0.2248(12)	-0.483(2)	0.030(2)	0.0193(14)	0.042(3)	0.023(3)	0.046(3)	0.029(3)
C18	0.279(1)	0.1298(10)	-0.445(2)	0.021(1)	0.0112(9)	0.059(3)	0.013(2)	0.039(3)	0.020(3)
C19	0.245(1)	0.1833(12)	-0.318(2)	0.021(1)	0.0138(13)	0.073(4)	0.007(2)	0.055(3)	0.020(3)
C20	0.238(1)	0.8195(10)	-0.322(2)	0.022(2)	0.0066(8)	0.076(5)	0.009(2)	0.013(3)	0.008(3)
C21	0.231(1)	0.4794(9)	0.383(2)	0.016(1)	0.0098(8)	0.046(3)	0.008(2)	0.033(4)	0.018(2)
C22	0.112(1)	0.4753(9)	0.485(2)	0.021(2)	0.0062(8)	0.054(4)	0.007(2)	0.015(4)	0.005(3)
C23	0.054(1)	0.4987(10)	0.118(2)	0.017(2)	0.0078(8)	0.072(5)	0.011(2)	0.028(4)	0.015(3)
C24	0.063(1)	0.3139(9)	0.092(2)	0.019(1)	0.0068(7)	0.053(3)	0.007(2)	0.026(3)	0.015(2)
C25	0.585(1)	0.2465(10)	-0.868(2)	0.028(2)	0.0099(9)	0.034(3)	-0.003(2)	0.021(3)	0.008(2)
C26	0.549(2)	0.1549(8)	-0.073(2)	0.036(2)	0.0050(6)	0.035(3)	-0.015(2)	0.023(4)	0.001(2)
C27	-0.5914(1)	0.8829(12)	-0.213(2)	0.015(1)	0.0159(12)	0.069(4)	0.001(2)	0.031(3)	0.030(3)
C28	0.5918(1)	0.8889(10)	-0.216(2)	0.027(2)	0.0145(9)	0.050(3)	0.002(2)	0.042(3)	0.023(2)
C29	0.482(1)	0.4947(9)	0.258(2)	0.020(1)	0.0125(9)	0.043(3)	0.016(2)	0.028(3)	0.023(2)
C30	0.586(2)	0.4878(13)	0.214(2)	0.037(2)	0.0202(12)	0.035(4)	0.042(2)	-0.007(5)	-0.014(4)
C31	0.472(2)	0.3397(10)	0.235(2)	0.050(3)	0.0077(8)	0.052(3)	-0.021(2)	0.050(4)	0.002(3)
C32	0.493(1)	0.2515(10)	0.189(2)	0.018(2)	0.0189(9)	0.037(3)	-0.003(2)	0.020(3)	0.004(3)
C33	0.449(1)	0.1855(9)	0.221(2)	0.018(2)	0.0063(7)	0.055(4)	0.007(2)	0.023(4)	0.009(3)
C34	0.451(1)	0.0937(9)	0.155(3)	0.017(2)	0.0044(7)	0.092(6)	0.002(2)	0.022(5)	0.009(3)
C35	0.393(1)	0.0288(10)	0.154(2)	0.014(1)	0.0079(9)	0.072(5)	0.006(2)	0.023(4)	0.006(3)
C36	0.269(1)	0.0201(9)	0.093(2)	0.020(2)	0.0054(7)	0.053(3)	0.001(2)	0.030(3)	0.009(3)

Table of Positional and Thermal Parameters and Their Estimated Standard Deviations (continued).

Atom	<i>x</i>	<i>y</i>	<i>z</i>	B.A.	Atom	<i>x</i>	<i>y</i>	<i>z</i>	B.A.
H1	5.2515(8)	5.6922(8)	5.3981(8)	5.0000(8)	H39	5.0000(8)	5.0000(8)	5.0000(8)	5.0000(8)
H2	5.2693(8)	5.8319(8)	5.4711(8)	5.0000(8)	H31	5.0000(8)	5.0000(8)	5.0000(8)	5.0000(8)
H3	5.3773(8)	5.9114(8)	5.4806(8)	5.0000(8)	H32	-5.0182(8)	5.0138(8)	5.0038(8)	5.0000(8)
H4	5.4536(8)	5.8363(8)	5.2248(8)	5.0000(8)	H33	5.0000(8)	5.0000(8)	5.0000(8)	5.0000(8)
H5	5.4482(8)	5.6858(8)	5.1419(8)	5.0000(8)	H34	5.0000(8)	5.0000(8)	5.0000(8)	5.0000(8)
H6	5.5579(8)	5.1851(8)	5.4823(8)	5.0000(8)	H35	5.0181(8)	5.0000(8)	-5.0151(8)	5.0000(8)
H7	5.5471(8)	5.3347(8)	5.5696(8)	5.0000(8)	H36	-5.0089(8)	5.2418(8)	-5.0067(8)	5.0000(8)
H8	5.1232(8)	5.4879(8)	5.3983(8)	5.0000(8)	H37	5.0000(8)	5.0000(8)	5.0181(8)	5.0000(8)
H9	5.2247(8)	5.3112(8)	5.1263(8)	5.0000(8)	H38	5.0000(8)	5.0000(8)	5.0000(8)	5.0000(8)
H10	5.2578(8)	5.1813(8)	5.0371(8)	5.0000(8)	H39	-5.0059(8)	5.0059(8)	-5.0049(8)	5.0000(8)
H11	5.2828(8)	5.5195(8)	5.1128(8)	5.0000(8)	H40	-5.0068(8)	5.0000(8)	5.0000(8)	5.0000(8)
H12	5.1813(8)	5.4749(8)	5.0358(8)	5.0000(8)	H41	-5.0012(8)	-5.0372(8)	-5.0235(8)	5.0000(8)
H13	5.2336(8)	5.3618(8)	5.1392(8)	5.0000(8)	H42	-5.0048(8)	-5.0158(8)	-5.0000(8)	5.0000(8)
H14	5.3392(8)	5.4537(8)	5.1377(8)	5.0000(8)	H43	5.0000(8)	5.0000(8)	5.0273(8)	5.0000(8)
H15	5.2742(8)	5.3935(8)	5.3894(8)	5.0000(8)	H44	5.0000(8)	5.0000(8)	5.0000(8)	5.0000(8)
H16	5.1581(8)	5.3887(8)	5.3945(8)	5.0000(8)	H45	5.0000(8)	5.0000(8)	5.0000(8)	5.0000(8)
H17	5.1731(8)	5.2581(8)	5.5398(8)	5.0000(8)	H46	5.0000(8)	5.0000(8)	5.0000(8)	5.0000(8)
H18	5.1637(8)	5.2497(8)	5.3782(8)	5.0000(8)	H47	5.0000(8)	5.0000(8)	5.0000(8)	5.0000(8)
H19	5.3375(8)	5.2468(8)	5.2894(8)	5.0000(8)	H48	5.0000(8)	5.0000(8)	5.0000(8)	5.0000(8)
H20	5.3312(8)	5.2385(8)	5.4663(8)	5.0000(8)	H49	5.0000(8)	5.0000(8)	5.0000(8)	5.0000(8)
H21	5.3448(8)	5.1895(8)	5.4444(8)	5.0000(8)	H50	5.0000(8)	5.0000(8)	5.0000(8)	5.0000(8)
H22	5.2255(8)	5.1847(8)	5.5588(8)	5.0000(8)	H51	5.0000(8)	5.0000(8)	5.0000(8)	5.0000(8)
H23	5.1769(8)	5.1196(8)	5.3291(8)	5.0000(8)	H52	5.0000(8)	5.0000(8)	5.0000(8)	5.0000(8)
H24	5.2958(8)	5.1361(8)	5.2124(8)	5.0000(8)	H53	5.0000(8)	5.0000(8)	5.0000(8)	5.0000(8)
H25	5.1941(8)	5.0135(8)	5.4318(8)	5.0000(8)	H54	5.0000(8)	5.0000(8)	5.0000(8)	5.0000(8)
H26	5.3088(8)	5.5554(8)	5.3084(8)	5.0000(8)	H55	5.0000(8)	5.0000(8)	5.0000(8)	5.0000(8)
H27	5.2579(8)	5.5168(8)	5.4148(8)	5.0000(8)	H56	5.0000(8)	5.0000(8)	5.0000(8)	5.0000(8)
H28	5.2448(8)	5.4217(8)	5.2979(8)	5.0000(8)	H57	5.0000(8)	5.0000(8)	5.0000(8)	5.0000(8)
H29	5.0953(8)	5.5282(8)	5.2233(8)	5.0000(8)	H58	5.0000(8)	5.0000(8)	5.0000(8)	5.0000(8)

The form of the anisotropic thermal parameter is:

$$\exp[-\alpha(1,1)*h^1 + \alpha(2,2)*h^2 + \alpha(3,3)*h^3 + \alpha(1,2)*hk + \alpha(1,3)*hl + \alpha(2,3)*kl]$$
.

Estimated standard deviations in the least significant digits are shown in parentheses

Table of Bond Distances in Angstroms

Atom1	Atom2	Distance	Atom1	Atom2	Distance	Atom1	Atom2	Distance
SN1	C1	2.162(11)	C7	C8	1.419(15)	C22	C23	1.35(2)
SN1	C13	2.11(2)	C7	C12	1.363(14)	C23	C24	1.508(15)
SN1	C21	2.141(14)	C8	C9	1.37(2)	C24	C25	1.475(15)
SN1	C29	2.161(13)	C9	C10	1.33(2)	C25	C26	1.57(2)
SN2	C7	2.115(18)	C10	C11	1.32(2)	C26	C27	1.41(2)
SN2	C20	2.13(2)	C11	C12	1.375(14)	C27	C28	1.41(2)
SN2	C28	2.133(15)	C13	C14	1.27(2)	C29	C30	1.44(2)
SN2	C36	2.136(14)	C14	C15	1.49(2)	C30	C31	1.26(3)
C1	C2	1.344(14)	C15	C16	1.51(2)	C31	C32	1.45(2)
C1	C6	1.408(15)	C16	C17	1.39(2)	C32	C33	1.46(2)
C2	C3	1.386(15)	C17	C18	1.46(2)	C33	C34	1.467(15)
C3	C4	1.31(2)	C18	C19	1.53(2)	C34	C35	1.28(2)
C4	C5	1.34(2)	C19	C20	1.34(2)	C35	C36	1.56(2)
C5	C6	1.39(2)	C21	C22	1.50(2)			

Numbers in parentheses are estimated standard deviations in the least significant digits.

DL/413/83/01
GEN/413-2TECHNICAL REPORT DISTRIBUTION LIST, GEN

<u>No.</u> <u>Copies</u>		<u>No.</u> <u>Copies</u>	
2	Office of Naval Research Attn: Code 413 800 N. Quincy Street Arlington, Virginia 22217	Naval Ocean Systems Center Attn: Technical Library San Diego, California 92152	1
1	ONR Pasadena Detachment Attn: Dr. R. J. Marcus 1030 East Green Street Pasadena, California 91106	Naval Weapons Center Attn: Dr. A. B. Amster Chemistry Division China Lake, California 93555	1
1	Commander, Naval Air Systems Command Attn: Code 310C (H. Rosenwasser) Washington, D.C. 20360	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
1	Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko Port Hueneme, California 93401	Dean William Tolles Naval Postgraduate School Monterey, California 93940	1
1	Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
12	Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	Mr. Vincent Schaper DTNSRDC Code 2830 Annapolis, Maryland 21402	1
1	DTNSRDC Attn: Dr. G. Bosmajian Applied Chemistry Division Annapolis, Maryland 21401	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	Mr. A. M. Anzalone Administrative Librarian PLASTEC/ARRADCOM Bldg 3401 Dover, New Jersey 07801	1

DL/413/83/01
053/413-2

TECHNICAL REPORT DISTRIBUTION LIST, 053

Dr. M. F. Hawthorne
Department of Chemistry
University of California
Los Angeles, California 90024

Dr. D. Venezky
Chemistry Division
Naval Research Laboratory
Washington, D.C. 20375

Professor O. T. Beachley
Department of Chemistry
State University of New York
Buffalo, New York 14214

Dr. A. Cowley
Department of Chemistry
University of Texas
Austin, Texas 78712

Dr. W. Hatfield
Department of Chemistry
University of North Carolina
Chapel Hill, North Carolina 27514

Professor Richard Eisenberg
Department of Chemistry
University of Rochester
Rochester, New York 14627

Professor K. Neidenzu
Department of Chemistry
University of Kentucky
Lexington, Kentucky 40506

Dr. T. Marks
Department of Chemistry
Northwestern University
Evanston, Illinois 60201

Dr. J. Zuckerman
Department of Chemistry
University of Oklahoma
Norman, Oklahoma 73019

Professor K. M. Nicholas
Department of Chemistry
Boston College
Chestnut Hill, Massachusetts 02167

Professor R. Neilson
Department of Chemistry
Texas Christian University
Fort Worth, Texas 76129

Professor M. Newcomb
Department of Chemistry
Texas A&M University
College Station, Texas 77843

Professor R. Wells
Department of Chemistry
Duke University
Durham, North Carolina 27706

Professor K. O. Christe
Rockwell International
Canoga Park, California 91304

SEARCHED

8